

## Progress Report

- Title:** Selective Control of Annual Bluegrass (*Poa annua* L.) in Creeping Bentgrass Putting Greens
- Investigators:** Jim Baird, Brent Barnes, Alea Miehl, and Vanessa Ferrel  
Department of Botany and Plant Sciences  
2137 Batchelor Hall  
University of California, Riverside  
Riverside, CA 92521  
951-827-5630 Office  
951-333-9052 Cell  
jbaird@ucr.edu
- Sponsors:** Golf Course Superintendents Association of Northern California  
Northern California Golf Association  
San Diego Golf Course Superintendents Association  
Sierra Nevada Golf Course Superintendents Association  
Southern California Golf Association  
Southern California Golf Course Superintendents Association  
Southern California Section, PGA
- Cooperators:** Gary Custis, PBI Gordon  
Doug Houseworth, Arysta LifeScience Corporation  
Suk-Jin Koo, Moghu Research Center  
Bob Mack, Helena Chemical  
Todd Mayhew, Valent Corporation  
Dean Mosdell, Syngenta  
Chris Olsen, Bayer Environmental Science
- Objectives:** Evaluate existing and experimental herbicides for selective removal of annual bluegrass that persists in creeping bentgrass putting green turf.
- Evaluate herbicides and herbicide combinations for potential bentgrass injury prior to inclusion in experiments on golf courses throughout the State.
- Location:** UCR Turfgrass Research Facility, Riverside CA
- Soil:** Loamy sand amended with sand topdressing
- Experimental Design:** Randomized complete block; three replications
- Plot Size:** 3 ft x 6 ft

**Species/Cultivars:** Creeping Bentgrass (*Agrostis stolonifera* L.) ‘Cobra’

**Mowing Height:** 0.180 inches; 3 days/wk

**Irrigation:** 80% ETo (historical from previous week)/DU

**Cultivation:** alternate bi-monthly verticutting/solid-tine aeration; topdressing monthly

**Sprayer:** Bicycle with two 8003VS nozzles  
20-inch spacing  
35 psi  
510 ml/nozzle/30 sec = 80 GPA  
Metronome = 50 beats = 1 mph

**Application Dates:** A = May 13 and June 10, 2010  
B = May 13, 20, 28 and June 4, 2010  
C = Bi-weekly beginning May 13, 2010  
D = June 10, 2010  
E = Scheduled for September 1, 2010

**Data Collected:** *Poa annua* control (0-100%) based on initial *Poa* cover in each plot  
Bentgrass quality (1-9, 1 = dead; 6 = minimally acceptable)

### Results:

- ✓ From May 13 to July 14, average minimum and maximum air temperatures were 59.1 F (45.7 F low) and 85.0 F (99.2 F high), respectively; average minimum and maximum relative humidity was 35% (8% low) and 80% (85% high), respectively; average soil temperatures ranged from 62.5 F to 76.0 F (73.3 F average); total CIMIS ETo was 2.78 inches; no precipitation.
- ✓ More than one application of amicarbazone applied at 2.0 oz/A or greater caused severe injury or death to bentgrass turf (Figure 1; Table 1). Preliminary results from studies in Northern California indicate that these rates can be safely applied in cooler climates or during cooler periods of the year, but they are too high for late spring/early summer in Riverside.
- ✓ Maximum air temperatures occurred between the July 6 and July 14 rating dates. Furthermore, the green was vertical mowed on July 7. The combination of both stresses resulted in the lowest overall bentgrass quality ratings on July 14, and plots treated with higher rates of HM 0814 began showing significant turf injury relative to the control.
- ✓ MRC-01 provided the best combination of *Poa* control and bentgrass safety during this phase of the experiment (Table 2); however, there appeared to a minimum total dosage required to achieve optimal control. Hence, the 3.0 oz/1,000 ft<sup>2</sup> rate (surfactant vs. no surfactant) was added on to the treatment list later during the experiment.

- ✓ The Riverside study will continue along with ongoing studies on golf courses Northern California, and new studies to be initiated on golf courses in Southern California. Focus will be on refining application rates and frequencies of MRC-01, evaluation of tank-mix partners with MRC-01 and with amicarbazone at 1.0 oz/A, and evaluation of higher rates and/or more frequent applications of other herbicides in this study to achieve maximum *Poa* control with minimal bentgrass injury.




Figure 1. Bentgrass injury on July 6 following four weekly applications of amicarbazone at 2 oz/A from May 13 to June 4, 2010. Riverside, CA.

Table 1. Creeping bentgrass quality (1-9, 1=dead, 6=minimally acceptable) following application of herbicides. Riverside, CA. 2010.

Trt	Rate	Timing	5/20	5/28	6/6	6/19	6/25	7/6	7/14
Velocity	4 oz/A	A	7.0	7.7	7.7	8.0	9.0	9.0	8.0
Velocity + Trimmit	2 oz/A + 8 oz/A	A A	7.0	8.0	8.0	8.0	9.0	9.0	8.0
Amicarbazone	1 oz/A	A	8.0	8.0	8.0	8.0	8.0	9.0	7.3
Amicarbazone	2 oz/A	A	7.7	7.7	8.0	6.7	6.0	8.3	8.0
Amicarbazone	4 oz/A	A	7.0	6.0	6.3	2.7	2.3	2.7	3.0
Amicarbazone + Trimmit	2 oz/A + 8 oz/A	A A	7.0	6.7	7.3	5.0	4.7	6.0	7.7
MRC-01	1.25 oz/1,000ft <sup>2</sup>	A	8.0	8.0	8.0	9.0	9.0	8.7	7.0
Prograss	8 oz/A	A	8.0	8.0	7.7	8.7	9.0	9.0	7.7
Prograss + Amicarbazone	6 oz/A + 2 oz/A	A A	8.0	8.0	8.0	7.7	7.0	8.7	7.7
HM 0814	3 oz/1,000ft <sup>2</sup>	A	8.0	8.0	8.0	8.3	8.7	8.0	6.7
HM 0814	6 oz/1,000ft <sup>2</sup>	A	8.0	8.0	8.0	8.3	8.0	8.0	6.7
HM 0814 + Trimmit	2 oz/1000ft <sup>2</sup> + 8 oz/A	A A	8.0	8.0	8.0	8.3	8.3	8.0	7.3
Trimmit	10 oz/A	A	8.0	8.0	8.0	8.0	8.3	9.0	8.3
Trimmit	16 oz/A	A	8.0	8.0	8.3	9.0	8.3	9.0	8.7
Bensumec 4 LF	9.4 oz/1000ft <sup>2</sup>	E	8.0	8.0	8.0	8.7	9.0	9.0	7.0
SureGuard	6 oz/A	E	8.0	8.0	8.0	9.0	9.0	9.0	7.7
Amicarbazone	1 oz/A	B	7.7	7.7	6.0	6.7	6.7	8.3	8.0
Amicarbazone	2 oz/A	B	7.3	6.0	2.0	1.7	1.3	2.7	3.7
Amicarbazone + Trimmit	1 oz/A + 2 oz/A	B B	8.0	7.7	5.0	6.3	6.7	8.0	8.0
FeSO <sub>4</sub> <sup>1</sup>	16 oz/1000ft <sup>2</sup>	C	9.0	8.0	8.0	9.0	9.0	9.0	9.0
MRC-01	0.75 oz/1000ft <sup>2</sup>	B	8.0	8.0	7.7	9.0	9.0	8.7	7.7
MRC-01	3 oz/1000ft <sup>2</sup>	D	8.0	8.0	8.0	8.7	8.7	9.0	7.3
MRC-01 <sup>2</sup>	3 oz/1000ft <sup>2</sup>	D	8.0	8.0	8.0	8.3	9.0	9.0	7.7
Untreated Control	-	-	8.0	8.0	8.0	9.0	9.0	9.0	8.0
<b>LSD (0.05)*</b>			<b>0.3</b>	<b>0.4</b>	<b>0.6</b>	<b>0.8</b>	<b>1.0</b>	<b>1.0</b>	<b>1.1</b>

\*Treatment mean differences in columns greater than or equal to LSD are significantly different, Fisher's Protected LSD,  $P=0.05$ .

<sup>1</sup>FeSO<sub>4</sub> applied in 320 GPA of water; all other treatments applied in 80 GPA. <sup>2</sup>No surfactant added; all other treatments applied with 0.25% non-ionic surfactant.

Table 2. Annual bluegrass control (0-100%) following application of herbicides. Riverside, CA. 2010.

Trt	Rate	Timing	5/28	6/6	6/19	6/25	7/6	7/14
Velocity	4 oz/A	A	28	33	48	57	70	65
Velocity + Trimmit	2 oz/A + 8 oz/A	A A	31	25	40	34	73	48
Amicarbazone	1 oz/A	A	29	44	57	60	47	64
Amicarbazone	2 oz/A	A	37	52	96	99	80	92
Amicarbazone	4 oz/A	A	70	87	100	100	100	100
Amicarbazone + Trimmit	2 oz/A + 8 oz/A	A A	40	71	100	100	94	89
MRC-01	1.25 oz/1,000ft <sup>2</sup>	A	20	61	84	75	82	87
Prograss	8 oz/A	A	12	32	52	44	36	37
Prograss + Amicarbazone	6 oz/A + 2 oz/A	A A	38	45	90	88	78	68
HM 0814	3 oz/1,000ft <sup>2</sup>	A	10	20	25	21	23	30
HM 0814	6 oz/1,000ft <sup>2</sup>	A	21	18	54	48	51	54
HM 0814 + Trimmit	2 oz/1,000ft <sup>2</sup> + 8 oz/A	A A	19	9	31	22	52	63
Trimmit	10 oz/A	A	10	14	24	19	47	33
Trimmit	16 oz/A	A	18	24	32	28	51	62
Bensumec 4 LF	9.4 oz/1,000ft <sup>2</sup>	E	4	18	51	30	39	54
SureGuard	6 oz/A	E	17	10	34	27	21	17
Amicarbazone	1 oz/A	B	32	84	87	78	61	74
Amicarbazone	2 oz/A	B	69	98	100	100	100	100
Amicarbazone + Trimmit	1 oz/A + 2 oz/A	B B	21	82	100	97	87	76
FeSO <sub>4</sub>	16 oz/1,000ft <sup>2</sup>	C	20	23	26	28	49	40
MRC-01	0.75 oz/1,000ft <sup>2</sup>	B	32	86	97	98	76	74
MRC-01	3 oz/1,000ft <sup>2</sup>	D	10	20	50	45	56	98
MRC-01 <sup>2</sup>	3 oz/1,000ft <sup>2</sup>	D	23	25	34	27	62	91
Untreated Control	-	-	17	11	22	19	11	30
<b>LSD (0.05)*</b>			<b>21</b>	<b>23</b>	<b>21</b>	<b>24</b>	<b>29</b>	<b>29</b>

\*Treatment mean differences in columns greater than or equal to LSD are significantly different, Fisher's Protected LSD,  $P=0.05$ .

<sup>1</sup>FeSO<sub>4</sub> applied in 320 GPA of water; all other treatments applied in 80 GPA. <sup>2</sup>No surfactant added; all other treatments applied with 0.25% non-ionic surfactant.